


# Stabilis





































## Amlodipine besylate



### Stabilité des préparations

		30 mg Norvasc®	Oral Mix® >> 30 ml	23-27°C		60			3893
30 mg ® = Medisca		Pharmaceutique Inc	Oral Mix® >> 30 ml	23-27°C		60			3893
		30 mg Norvasc®	Oral Mix® >> 30 ml	3-7°C		90			3893
30 mg ® = Medisca		Pharmaceutique Inc	Oral Mix® >> 30 ml	3-7°C		90			3893
		69 mg ® = ?	POH benzoate de methyle 15% 287 mg Sirop simple 32 g Eau purifiée >> 100 ml	4°C		365			3886
		150 mg ® = ?	SyrSpend SF® PH 4 >> 150 mL	2-8°C		90			3819
		150 mg ® = ?	SyrSpend SF® PH 4 >> 150 mL	20-25°C		90			3819

		25 mg Norvasc®	Methylcellulose 125 mg POH benzoate de methyle 2,5 mg POH benzoate de propyle 1,25 mg Eau purifiée >> 12,5 ml Sirop simple >> 25 ml	25°C		56			2689
		25 mg Norvasc®	OraPlus® / OraSweet® (1:1) >> 25 ml	25°C		56			2689
		25 mg Norvasc®	Methylcellulose 125 mg POH benzoate de methyle 2,5 mg POH benzoate de propyle 1,25 mg Eau purifiée >> 12,5 ml Sirop simple >> 25 ml	4°C		90			2689
		25 mg Norvasc®	OraPlus® / OraSweet® (1:1) >> 25 ml	4°C		90			2689
		69 mg ® = ?	POH benzoate de methyle 15% 287 mg Sirop simple 32 g Eau purifiée >> 100 ml	4°C		365			3886
		1 mg Norvasc®	Oral Mix® >> 1 ml	23-27°C		45			3893
1 mg ® = Medisca Pharmaceutique Inc			Oral Mix® >> 1 ml	23-27°C		45			3893
		1 mg Norvasc®	Oral Mix® >> 1 ml	3-7°C		90			3893
1 mg ® = Medisca Pharmaceutique Inc			Oral Mix® >> 1 ml	3-7°C		90			3893




## Bibliographie

	Type	Source
2689	Revue	Nahata MC, Morosco RS, Hipple TF. Stability of amlodipine mesylate in two liquid dosage form. J Am Pharm Assoc 1999 ; 39: 375-377.
3819	Revue	Ferreiraa A.O, Poloninia H.C, Silva S.L, Patricio F.B, Brand?oa M.A.F, Raposoa N.R.B. Feasibility of amlodipine besylate, chloroquine phosphate, dapson, phenytoin, pyridoxine hydrochloride, sulfadiazine, sulfasalazine, tetracycline hydrochloride, trimethoprim and zonisamide in SyrSpend® SF PH4 oral suspensions. J Pharm Biomed Anal 2015 ; 118: 105-112.
3886	Revue	van der Vossen A.C, van der Velde I, Smeets O.S.N.M., Postma D.J, Vermes A., Koch B.C.P, Vulto A.G., Hanff L.M. Design and stability study of an oral solution of amlodipine besylate for pediatric patients. Eur J Pharm Sci 2016 ; 92 : 220-223.
3893	Revue	Friciu M, Zaraa S, Leclair G. Stability of Extemporaneously Compounded Amlodipine Besylate Oral Suspensions Can J Hosp Pharm 2016 ; 69, 4: 327-329.



## Dictionnaire

 Antihypertenseur	 Solution buvable
 Stabilité des préparations	 Contenant
 Origine	 Excipient
 Température	 Conservation
 Durée de stabilité	 Biosimilaire
 Données conflictuelles	 Bibliographie
 Verre	 Comprimés
 A l'abri de la lumière	 Jour
 Poudre	 Flacon plastique
 Lumière	 Seringue PP orale
 Bibliographie	 Dictionnaire