


Stabilis



Ursodesoxycholic acid



Stabilité des préparations

| | | 5000 mg ® = ? | Inorpha® 100 mL Gomme xanthane 0,25% 250 mg | 2-6°C | | 90 | | | 4498 |
|--|--|-------------------|--|---------|--|-----|--|--|------|
| | | 5000 mg ® = ? | Inorpha® 100 mL Gomme xanthane 0,25% 250 mg | 20-24°C | | 60 | | | 4498 |
| | | 3600 mg Actigal® | Glycérine Sirop simple >> 60 ml | 4°C | | 35 | | | 2449 |
| | | 3000 mg ® = ? | SyrSpend SF PH4® >> 150 ml | 2-8°C | | 90 | | | 4177 |
| | | 3000 mg Urso® | OraSweet SF® / sirop de fraise (1:1) | 2-5°C | | 90 | | | 2410 |
| | | 3000 mg Urso® | OraPlus® / OraSweet SF® (1:1) | 2-5°C | | 90 | | | 2410 |
| | | 4000 mg Ursodiol® | Acésulfame potassique 400 mg Glycosides de stéviol 95% 400 mg | 25°C | | 181 | | | 4214 |
| | | 8000 mg Ursodiol® | Acésulfame potassique 400 mg Glycosides de stéviol 95% 400 mg | 25°C | | 181 | | | 4214 |
| | | 3000 mg Urso® | OraSweet SF® / sirop de fraise (1:1) | 2-5°C | | 90 | | | 2410 |
| | | 3000 mg Urso® | OraPlus® / OraSweet SF® (1:1) | 2-5°C | | 90 | | | 2410 |
| | | 8000 mg Ursodiol® | Acésulfame potassique 400 mg Glycosides de stéviol 95% 400 mg | 5°C | | 181 | | | 4214 |
| | | 4000 mg Ursodiol® | Acésulfame potassique 400 mg Glycosides de stéviol 95% 400 mg | 5°C | | 181 | | | 4214 |

| | | | | | | | | | |
|---|--|-------------|--|--|----|--|--|--|------|
| ? | | 1,5 g ® = ? | Glycérine 20 g Methylcellulose 1% >> 100 mL 25°C PH= 4.4 | | 60 | | | | 3782 |
| ? | | 1,5 g ® = ? | Glycérine 20 g Methylcellulose 1% >> 100 mL 40°C PH= 4.4 | | 60 | | | | 3782 |
| ? | | 1,5 g ® = ? | Glycérine 20 g Methylcellulose 1% >> 100 mL 5°C PH= 4.4 | | 30 | | | | 3782 |



Bibliographie

| | Type | Source |
|------|-------|---|
| 2410 | Revue | Johnson CE, Streetman DD. Stability of oral suspensions of ursodiol made from tablets Am J Health-Syst Pharm 2002 ; 59: 361-363. |
| 2449 | Revue | Johnson CE, Nesbitt J. Stability of ursodiol in an extemporaneously compounded oral liquid Am J Health-Syst Pharm 1995 ; 52: 1798-1800. |
| 3782 | Revue | A. Santove?a, E. S?nchez-Negr?n, L. Charola, M. Llabr?s, J.B. Fari?a □ Study of quality and stability of ursodeoxycholic acid formulations for oral pediatric administration Int J Pharm 2014 , 477, 1-2 : 32-38. |
| 4177 | Revue | Polonini H, Loures da Silva S, Fernandes Brandao M.A, Bauters T, De Moerloose B, De Oliveira Ferreira A. Compatibility of Baclofen, Carvedilol, Hydrochlorothiazide, Mercaptopurine, Methadone Hydrochloride, Oseltamivir Phosphate, Phenobarbital, Propranolol Hydrochloride, Pyrazinamide, Sotalol Hydrochloride, Spironolactone, Tacrolimus Monohydrate, Ursodeoxycholic Acid, and Vancomycin Hydrochloride Oral Suspensions Compounded with SyrSpend SF PH4. Int J Pharm Compound 2018 ;22,6:516-526 |
| 4214 | Revue | Pramar Y.V, Mandal T.K, Bostanian L.A, Nguyen A.TQ, Miller V, Morris T.C, Graves R.A. Stability of Compounded Ursodiol Suspensions in PCCA Base, SuspendIt. Int J Pharm Compound 2019 ;23,1:70-76 |
| 4498 | Revue | Hausherr A, Roessle C, Pinet E, Vasseur V, Abarou T, Benakouche S, Bourdon O, Storme T. Development and validation of a new HPLC method for the analysis of a novel oral suspension formulation of 50 mg/mL ursodesoxycholic acid for newborns. Pharmaceutical Technology in Hospital Pharmacy 2020 |



Dictionnaire

| | | | |
|--|----------------------------|--|------------------|
| | Divers | | Solution buvable |
| | Stabilité des préparations | | Contenant |
| | Origine | | Excipient |
| | Température | | Conservation |
| | Durée de stabilité | | Biosimilaire |
| | Données conflictuelles | | Bibliographie |
| | Verre | | Poudre |
| | A l'abri de la lumière | | Jour |
| | Gélules | | Flacon plastique |
| | Non précisée | | Comprimés |
| | Non précisé | | Bibliographie |
| | Dictionnaire | | |